



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

CALCULUS.

101. Proposed by WILLIAM FRED FLEMING, Denison, Tex.

A 24-inch joint of 6-inch stove pipe is compressed at one end to make it fit over an elliptical opening in a stove (for the escape of smoke). The ellipse has a major axis of 8 inches. What reduction is there in the solid contents of the stove pipe, assuming that its compressed shape may be generated by a 6-inch circle which passes uniformly from one end to the other and perpendicular to the axis of the pipe ?

I. Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

The perimeter of an ellipse semi-major axis a is

$$4a \int_0^{\frac{1}{2}\pi} \sqrt{1-e^2 \sin^2 \theta} d\theta = s.$$

In this problem $a=4$ and $s=6\pi$, the circumference of the pipe.

$$\begin{aligned} \therefore 6\pi &= 16 \int_0^{\frac{1}{2}\pi} (1 - \frac{1}{2}e^2 \sin^2 \theta - \frac{1}{8}e^4 \sin^4 \theta - \frac{1}{16}e^6 \sin^6 \theta - \text{etc.}) d\theta \\ &= 8\pi (1 - \frac{1}{4}e^2 - \frac{3}{64}e^4 - \frac{5}{256}e^6 - \frac{175}{16384}e^8 - \frac{441}{65536}e^{10} - \text{etc.}) \end{aligned}$$

Let $e=u$. Then $u + \frac{3}{16}u^2 + \frac{5}{64}u^3 + \frac{175}{4096}u^4 + \frac{441}{16384}u^5 + \dots = 1$.

By reversion of series $u=e^2=.8013$.

$\therefore b=a\sqrt{1-e^2}=4\sqrt{.1987}=1.783$ inches.

Let the pipe be compressed uniformly for the entire length. Also let πxy be the area of any elliptical section, and z the distance of this section from the circular end of the pipe.

$$\text{Then } x = \frac{72+z}{24}, \quad y = \frac{72-1.217z}{24}.$$

$$V = \pi \int_0^{24} xy dz = \frac{\pi}{576} \int_0^{24} (5184 - 15.624z - 1.217z^2) dz.$$

$\therefore V=624.4568$ cubic inches.

V_1 = volume before compression.

$\therefore V_1 = 9\pi \times 24 = 678.5856$ cubic inches.

$\therefore V_1 - V = 54.1288$ cubic inches reduction in volume.

II. Solution by H. C. WHITAKER, Ph. D., Professor of Mathematics, Manual Training School, Philadelphia, Pa.

Denote the length of the pipe ($=24$ inches) by h , the semi-major axis ($=4$) of the ellipse by a , the semi-minor axis by b , the eccentricity by e , the radius of the circle ($=3$) by r . Then equating semi-perimeters :

$$\pi r = \pi a \left(1 - \frac{e^2}{2^2} - \frac{1^2 \cdot 3 \cdot e^4}{2^2 \cdot 4^2} - \frac{1^2 \cdot 3^2 \cdot 5 \cdot e^6}{2^2 \cdot 4^2 \cdot 6^2} - \right).$$

Solving by trial, $e = .895$, and $b = 1.784$.

At a distance of z from the circular end of the pipe, the section is an ellipse, the semi-axes of which are found to be

$$\frac{hr - rz + az}{h} \quad \text{and} \quad \frac{hr - rz + bz}{h}.$$

Hence the volume is

$$\int_0^h \pi \left(\frac{hr - rz + az}{h} \right) \left(\frac{hr - rz + bz}{h} \right) dz = \frac{\pi h}{6} (ar + br + 2r^2 + 2ab) = 198.5\pi.$$

Hence the loss $= 216\pi - 198.5\pi = 17.5\pi = 55$ cubic inches.

MISCELLANEOUS.

82. Proposed by A. H. BELL, Hillsboro, Ill.

Four spheres of equal radii $r=5$, are in contact, and form a triangular pyramid. How large is the sphere that can be placed in the middle and be in contact with the four spheres.

Solution by J. W. YOUNG, Fellow in Mathematics, Cornell University, Ithaca, N. Y., and J. SCHEFFER, A. M., Hagerstown, Md.

Let A, B, C, B' (Fig. 1) be the centers of the four spheres. They evidently form the corners of a regular tetrahedron. Fig. 2 is a picture of a plane section of the pyramid of spheres, passed through the points $AB'L$, where L is the point of tangency of the two spheres (C, B) .

From Fig. 1,

$$\left. \begin{aligned} AN/AM &= \sin 60^\circ \\ AN/AB' &= \cos 60^\circ \end{aligned} \right\}.$$

$$\therefore AB'/AM = \tan 60^\circ = \sqrt{3} = \sec \angle DAM.$$

In Fig. 2, then, $\angle DAM$ is $\sec^{-1} \sqrt{3}$. It is clear that the required small sphere must have its center on DM and must touch both spheres (A, D) . Let $\angle ADM = \theta$.

$$\text{Then } \sin \theta = 1/\sqrt{3}, \cos \theta = \sqrt{2}/3, DT/r = \sqrt{2}/3.$$

$$\therefore DT = (r/2) \sqrt{6}.$$

$$\therefore RT = DT - r = (r/2)(\sqrt{6} - 2) = \text{radius of small sphere.}$$

$$r=5 \text{ gives } RT = 1.1238.$$




Fig. 1.

Fig. 2.

85. Proposed by COOPER D. SCHMITT, A. M., Professor of Mathematics, University of Tennessee, Knoxville, Tenn.

Prove that at least one of the three sides of a rational right triangle must be divisible by 5.